

IN THE CLAIMS:

Please amend the claims as follows:

1. (Currently amended) A method of treating or preventing decreased nitric oxide formation resulting from sub-optimal urea cycle function in a subject, the method comprising:
 - (a) providing a subject under conditions of sub-optimal urea cycle function; and
 - (b) administering to ~~[[a]]~~ the subject ~~in need thereof~~ a therapeutically effective amount of a nitric oxide precursor, whereby treatment or prevention of decreased nitric oxide formation resulting from sub-optimal urea cycle function is accomplished.
2. (Original) The method of claim 1 wherein the administering is intravenously or orally.
3. (Original) The method of claim 1, wherein the sub-optimal urea cycle function further comprises decreased urea cycle intermediate production.
4. (Original) The method of claim 1, wherein the subject is suffering from a disorder associated with decreased urea cycle intermediate production or wherein the subject is exposed or about to be exposed to an environmental stimulus associated with decreased urea cycle intermediate production.
5. (Currently amended) The method of claim 4, wherein the disorder is selected from the group consisting of hepatitis, cirrhosis, pulmonary hypertension, necrotizing enterocolitis (NEC), Acute Respiratory Distress Syndrome, ethnic specific endothelial dysfunction, erectile dysfunction, ~~bone marrow transplant toxicity in a subject undergoing bone marrow transplant,~~ sepsis, asthma, and combinations thereof.
6. (Currently amended) The method of claim 4, wherein the environmental stimulus is selected from the group consisting of chemotherapy, cardiac surgery, increased oxidative stress, ~~bone marrow transplant,~~ septic shock, acute asthma attack, hypoxia, hepatotoxin exposure and combinations thereof.

7. (Original) The method of claim 1, wherein the nitric oxide precursor is selected from the group consisting of citrulline, arginine and combinations thereof.
8. (Original) The method of claim 1, wherein the nitric oxide precursor is administered in a dose ranging from about 100 mg to about 30,000 mg.
9. (Original) The method of claim 8, wherein the nitric oxide precursor is administered in a dose ranging from about 250 mg to about 1,000 mg.
10. (Original) The method of claim 1, wherein the subject is a human.
- 11-17. (Canceled)
18. (Currently amended) A method of treating or preventing ~~a disorder selected from the group consisting hepatitis, cirrhosis, pulmonary hypertension, necrotizing enterocolitis (NEC), Acute Respiratory Distress Syndrome, ethnic specific endothelial dysfunction, erectile dysfunction, asthma, and combinations thereof~~ in a subject suffering from sub-optimal urea cycle function, the method comprising administering to a subject in need thereof a therapeutically effective amount of a nitric oxide precursor, whereby the pulmonary hypertension is treated or prevented.
19. (Original) The method of claim 18, wherein the administering is intravenously or orally.
20. (Original) The method of claim 18, wherein the nitric oxide precursor is selected from the group consisting of citrulline, arginine and combinations thereof.
21. (Original) The method of claim 18, wherein the nitric oxide precursor is administered in a dose ranging from about 100 mg to about 30,000 mg.
22. (Original) The method of claim 21, wherein the nitric oxide precursor is administered in a dose ranging from about 250 mg to about 1,000 mg.
23. (Original) The method of claim 18, wherein the subject is a human.
24. (Canceled)
25. (Currently amended) A method of raising a level of a nitric ~~[[acid]]~~ oxide precursor in a subject ~~in need thereof~~ suffering from sub-optimal urea cycle function, the method comprising administering to the subject a therapeutically

effective amount of a nitric oxide precursor, whereby a level of a nitric oxide precursor in the subject is raised.

26. (Original) The method of claim 25, wherein the administering is intravenously or orally.
27. (Original) The method of claim 25, wherein the nitric oxide precursor is selected from the group consisting of citrulline, arginine and combinations thereof.
28. (Original) The method of claim 25, wherein the nitric oxide precursor is administered in a dose ranging from about 100 mg to about 30,000 mg.
29. (Original) The method of claim 28, wherein the nitric oxide precursor is administered in a dose ranging from about 250 mg to about 1,000 mg.
- 30-33. (Canceled)
34. (Previously presented) The method of claim 6, wherein the environmental stimulus comprises increased postoperative pulmonary vascular tone associated with cardiac surgery.

Please add the following new claims:

35. (New) A method of treating or preventing decreased nitric oxide formation resulting from sub-optimal urea cycle function in a subject suffering from a disorder associated with decreased urea cycle intermediate production, the method comprising:
 - (a) providing a subject suffering from a disorder associated with decreased urea cycle intermediate production, wherein the disorder is pulmonary hypertension; and
 - (b) administering to the subject a therapeutically effective amount of citrulline, whereby treatment or prevention of decreased nitric oxide formation resulting from sub-optimal urea cycle function is accomplished.
36. (New) A method of treating or preventing decreased nitric oxide formation resulting from sub-optimal urea cycle function in a subject exposed to an environmental stimulus associated with decreased urea cycle intermediate production, the method comprising:

- (a) providing a subject exposed to an environmental stimulus associated with decreased urea cycle intermediate production, wherein the environmental stimulus is cardiac surgery; and
 - (b) administering to the subject a therapeutically effective amount of citrulline, whereby treatment or prevention of decreased nitric oxide formation resulting from sub-optimal urea cycle function is accomplished.
- 37. (New) A method of treating or preventing pulmonary hypertension in a subject suffering from sub-optimal urea cycle function, the method comprising administering to a subject in need thereof a therapeutically effective amount of citrulline, whereby the pulmonary hypertension is treated or prevented.
- 38. (New) A method of raising a level of a nitric oxide precursor in a subject suffering from sub-optimal urea cycle function, the method comprising administering to the subject a therapeutically effective amount of citrulline, whereby the level of a nitric oxide precursor in the subject is raised.